

MELATONIN
Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

DOCUMENT TO BE SCANNED INTO ELECTRONIC RECORDS AND FILED IN NOTES

Patient name:	Date of Birth:	NHS number:
Patient Address:	Referring Consultant/Paediatrician name and contact details:	GP name and contact details:

INTRODUCTION

Melatonin is a pineal hormone which affects sleep. It is normally secreted at night and its main function is the regulation of circadian rhythm and sleep. It plays an important role, in setting the correct timing of sleep - wake cycles.

The administration of exogenous melatonin has a rapid, transient, mild sleep inducing effect and it lowers alertness, body temperature and performance for about 3 to 4 hours after the administration of low doses of immediate release formulations.

Clinical experience suggests that it may be of value for treating sleep onset insomnia and delayed sleep phase syndrome in children with conditions such as visual impairment, cerebral palsy, attention deficit hyperactivity disorder, autism spectrum disorder, and learning difficulties. In practice, the use of melatonin for the treatment of paediatric sleep-wake cycle disorders is widespread. There are a number of published trials, although these are often small and of short duration. As such it is difficult to draw firm conclusions.

Children with and or without co-morbid ADHD treated with melatonin have been shown to fall asleep earlier and sleep for longer when compared to controls. Generally no significant change in behaviour or attention have been demonstrated. It would appear that there is wide variability in response. Melatonin may be most effective in those children whose sleep patterns indicate that their circadian rhythm is disrupted, and in whom sleep hygiene methods have been ineffective.

Circadin® 2mg tablets are a sustained release formulation of Melatonin which is licensed in the UK as monotherapy for the treatment of primary insomnia in adults aged 55 years and over. Melatonin is not licensed for use in children. However its use off label is preferable to the use of an unlicensed special. The MHRA recommends that a licensed product should be used, even if it is used off label, before considering the use of an unlicensed preparation.

For the purpose of this guidance, the term 'specialist' is used to reference to a consultant paediatrician and child and adolescent mental health (CAMHS) consultant.

Please note: To take into account local differences in clinical practice between Luton/Bedfordshire and the London localities, certain statements apply to specific areas only as outlined in the documents. Luton/Bedfordshire are highlighted in blue and London localities highlighted in green.

MELATONIN
 Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

Patient group

Diagnosis of a sleep disorder in children made by a specialist in paediatrics, CAMHS (ELFT) and Learning Disabilities (LD).

For use in children of at least 1 year of age with neurodevelopment disability, autism spectrum disorder, visual impairment or neuropsychiatric disorders and chronic sleep disturbance, including chronic fatigue syndrome, where:

- Symptoms of sleep disturbance have been present for at least six months or sleep disturbance is so severe that it is causing significant family disturbance
- And after failure of sleep hygiene / behavioural measures.

CARE PATHWAY AND RESPONSIBILITY ARRANGEMENTS ACROSS PRIMARY AND SECONDARY CARE

Specialty	Indication	Duration	Prescribed by	Reviewed by
<i>For recommendation by a specialist and continuation by General Practitioner (GP)</i>				
Paediatric and neurology/ CAMHS (ELFT)/ LD	Insomnia in the above group	Duration of therapy to be agreed by specialist.	To be initiated by specialist and continuation by GP.	Annual review by the Paediatrician/ neurologist /CAMHS (ELFT)/LD team. 6 monthly review by GP in between annual specialist review.

Patient recommendation/transfer will be done via a letter from the Paediatric Consultant/CAMHS consultant/ LD consultant to the GP.

DOSE AND ADMINISTRATION

Age	Oral dose	Maximum dose
1 year – 18 years	Initially 2 mg daily. Increase every 1-2 weeks based on response to 4-6mg daily.	Usually 10mg daily
<ul style="list-style-type: none"> • Take the Melatonin after food. • Take dose 30-60 minutes before bedtime. • For children waking during the night, the same dose or a smaller dose can be repeated during the night (as prescribed by the specialist- to be communicated to the GP in writing). • Melatonin 2mg MR tablet (Circadin®) can be halved using a tablet cutter and it will retain its slow release characteristics. • For children <u>with tubes/those with severe swallowing difficulties</u> where <u>all medicines are liquids</u>, a liquid preparation of Melatonin is available (see third line under prescription and supply) • NOTE: crushing the MR tablet will mean that it is no longer modified release. (None of the currently available special formulations are slow release). • Refer to current BNFc and Summary of Product Characteristics for further information. 		

Please specify the brand of melatonin to be prescribed. This should be advised by the hospital.

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

PRESCRIPTION AND SUPPLY

Melatonin should be prescribed generically as Melatonin 2mg MR tablet. The brand Circadin® will be supplied as this is currently the only licensed product available.

Melatonin 2mg MR tablets should be labeled with advice to swallow the tablets whole and to avoid crushing.

Patients with difficulty swallowing, who will need to crush the tablets before administering, must be advised to override this instruction. The prescription should state that the medication is to be crushed prior to administration.

First line	Melatonin 2mg m/r tablets (Circadin®). (Cost 2017: £15.39 for 30 tablets). Circadin® tablets should be used first line if the appropriate dose can be obtained by either swallowing whole or crushing into water/ soft food.
Second Line	Melatonin 2mg m/r tablets (Circadin®). Can be halved without losing modified release characteristics. It can be crushed and mixed with water/juice/soft food for swallowing difficulties. Please note: once crushed it will no longer be modified release. London localities only: Biomelatonin 3mg tablets. If doses are for example, 3mg, 6mg or 9mg daily
Third line	Melatonin oral solution (Not the Melatonin suspension) is available as an unlicensed special order medicine. (Cost 07/2017: 5mg/5ml (200ml) for £52.70; 1mg/1ml sugar free (200ml) for £21.00) The specialist prescriber is advised to review use annually in the case of the oral solution. For children <u>with tubes/those with severe swallowing difficulties</u> where all medicines are liquids Please note: The oral solution should only be used in exceptional circumstances.

London localities only: Biomelatonin 3mg tablets. There has been feedback from the London areas that CCGs/GPs are happy to prescribe this formulation.

Refer to current BNFc and Summary of Product Characteristics for further information.

ADVERSE EFFECTS

Melatonin is generally well tolerated and adverse reactions reported are at similar levels to those reported with placebo. The long term side effects have not been evaluated.

Tiredness, headaches, dizziness, pharyngitis, abdominal back pain, constipation, dry mouth, weight gain, asthenia and irritability, sleep disorders, restlessness, nervousness and sweating have been reported following its use. Other rare side effects include restlessness, confusion, increased heart rate, itching and nausea.

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

Overdose

Administration of daily doses of up to 300mg of Melatonin without causing clinically significant adverse reactions have been reported in literature. High dose can reduce body temperature and if overdose occurs, drowsiness is to be expected. Clearance of the active substance is expected within 12 hours after ingestion. No special treatment is required.

Discontinuation does not appear to be associated with withdrawal effects. If any persistent and problematic side effects occur discontinue melatonin and refer the patient back to the hospital team

CAUTIONS

Cautions

- Drowsiness - driving or other activities that put the patient or others at risk should be avoided if the patient is affected by drowsiness
- Patients with rare hereditary problems of galactose intolerance, the LAPP lactase deficiency or glucose- galactose malabsorption should not take Circadin® brand melatonin (contains lactose)
- Patients with renal impairment
- Melatonin may worsen restless legs syndrome
- Melatonin can affect serotonin levels and should be used with caution in conjunction with other serotonin related agents.

Increased seizure activity has been reported in patients with epilepsy but there is also anecdotal evidence that seizure activity improves as a result of improved sleep. Much of the clinical trial data with Melatonin does not appear to report an increase in seizure frequency. However, the current data should be treated with caution due to the short term nature, size and heterogeneity of the populations studied. Prescribers are advised to use Melatonin with caution in young persons with epilepsy and to monitor for changes in seizure activity.

Endogenous serum Melatonin concentration is elevated in nocturnal asthmatic patients. Although the clinical trial data presented here does not indicate an increase in asthma symptoms, Melatonin should be used with caution in this group. Most commercial Melatonin is synthesized in the laboratory. However, in rare cases it has been derived from animal pineal glands. Melatonin from animal sources should be avoided due to the possibility of contamination.

Pregnancy and Breast Feeding

There are no known concerns with Melatonin in pregnancy. Where young girls or women taking Melatonin become pregnant, cases should be managed on an individual basis, seeking expert advice from the specialists.

Use in pregnant women and by women intending to become pregnant is not recommended. Breastfeeding is not recommended for women taking Melatonin.

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

CONTRAINDICATIONS AND INTERACTIONS

Contraindications

- Hypersensitivity to Melatonin or any excipients
- Patients with hepatic impairment
- Patients with autoimmune diseases or taking immunosuppressants

Drug interactions

Please refer to the most up-to date BNF and SPC for Melatonin for detailed information on interactions.

5 and 8-methoxypsoralen	Use with caution with melatonin, increased melatonin plasma concentration
Alcohol	Avoid – reduces effectiveness of melatonin
Carbamazepine and rifampicin	May give rise to reduced plasma concentrations of melatonin.
Cigarette smoking	May reduce melatonin concentrations
Cimetidine	Use with caution with melatonin, increased melatonin plasma concentration
Ciprofloxacin and other quinolones	Use with caution with melatonin, increased melatonin plasma concentration
Fluvoxamine	Avoid use with melatonin as melatonin plasma concentrations are markedly increased (metabolism inhibited)
Nifedipine	May affect blood pressure control
Oestrogens	Use with caution with melatonin, increased melatonin plasma concentration
Other hypnotics and CNS depressants	Melatonin may enhance the sedative properties of other drugs acting on the CNS e.g. benzodiazepines
Warfarin	INR may be increased

MONITORING AND REVIEW

No baseline monitoring is required. An annual review of Melatonin therapy should be completed by the specialist with the patient. In addition to this, the GP will review at the 6 month point between the specialist annual review. If possible, the patient and/ or carer should keep a sleep diary a month prior to the appointment which can form part of the review.

The specialist should consider a Melatonin free trial at least annually to assess the continued need for treatment. The outcome of any treatment break must be recorded in the patient's notes.

It is unusual for a child to continue melatonin into adulthood. It is the responsibility of the referring specialist to decide if it is appropriate to continue for a specific individual. The specialist will then advise the GP on future management.

Treatment should be stopped when there is a lack of effect based on information from the sleep diary, Melatonin break and patient/ parent perception.

The GP should be informed of any changes to Melatonin therapy.

ACTION AND ADVICE

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

If adverse effects are not tolerated, melatonin can be stopped immediately.

The patient's further therapy should be discussed with the patient's specialist.

SHARED CARE

Shared care guideline: is a document which provides information allowing patients to be managed safely by primary care, secondary care and across the interface. It assumes a partnership and an agreement between a hospital specialist, GP and the patient and also sets out responsibilities for each party. The intention to share care should be explained to the patient and accepted by them. Patients are under regular follow-up and this provides an opportunity to discuss drug therapy. Intrinsic in the shared care agreement is that the prescribing doctor should be appropriately supported by a system of communication and cooperation in the management of patients. The doctor who prescribes the medicine has the clinical responsibility for the drug and the consequence of its use.

Monitoring instructions and responsibilities

Specialist

- The specialist will recommend, initiate and prescribe the initial supply of 28 days of Melatonin. This will allow time for the GP to receive the information and accept shared care agreement.
- **Luton and Bedfordshire only: The Specialist will initiate treatment and prescribe until the patient is stabilised and the GP agrees to share care (minimum prescription supply is 28 days).'**
- Ensure the patient/ carer understands their treatment regimen and any monitoring or follow up that is required (using advocacy if appropriate).
- Ensure Melatonin treatment is reviewed annually by a member of the specialist team to monitor response, and to stop if there is lack of efficacy.
- A Melatonin free period should be agreed with the young person/ parent/ carer and the specialist to establish if the patient requires continued Melatonin treatment.
- Any Melatonin free period should be documented and reviewed at an appropriate interval (follow up) as agreed between the specialist/ young person/ parent/ carer.
- Ensure the patient/carers are aware of the off-license status or use of unlicensed preparation.
- Discuss benefits and side effects of treatment with the patient/ carer
- Document consent in the patient notes.
- Clinical supervision of the patient by routine clinic follow-up on a regular basis.
- Specialist should indicate, where relevant, in prescribing instructions that the tablet should be crushed.
- The specialist must supply the GP with a summary of the outpatient appointment within 21 days of initiation of treatment, where there are any changes, and where the Melatonin is discontinued.
- The letter to the GP should ensure the correct strength, dose and frequency are stated. This includes any specific instructions on e.g. frequency- repeated dose during the night, if the child wakes up; formulation- crushed or specific administration of liquid for NG.
- To ensure any correspondence to the GP includes the specialist contact details.
- The specialist will provide the GP with reference guidelines for the patient and ensure the patient/carers understand they will need to make an appointment to receive a

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

prescription/further supplies from the GP.

- Inform the GP if a non-standard dose is used and/or the preparation is changed from the agreed choice.
- Evaluate any reported adverse effects by GP or patient.
- Advise GP on review, duration or discontinuation of treatment where necessary.
- Inform GP of patients who do not attend clinic appointments.
- Ensure that backup advice and support is available for patient and GP at all times.
- Review the need for on-going Melatonin treatment for children reaching 18 years of age.
- Advise the GP of the arrangements for the future monitoring of the patient, should the young person need to continue melatonin once they reach adulthood.
- Report adverse events to the MHRA (yellow card reporting scheme) and inform GP.
- Where the off-label (Circadin®) preparation is not appropriate, provide clinical justification for using the alternative or liquid formulation in any correspondence sent to the GP.

General Practitioner

- Ensure that the patient/carer understands the nature, effect and potential side effects of the drug before prescribing it as part of the shared care programme and contact the specialist for clarification where appropriate.
- Prescribe Melatonin after communication with the specialist.
- Prescribers should indicate, if relevant as per communication from the specialist, in prescribing instructions that the tablet should be crushed.
- Monitor patient's health and wellbeing, if relevant as advised by specialist, in between specialist review appointments of physical health and well-being.
- Ask parent/ carer about effectiveness.
- Report any changes in physical health/ wellbeing and/ or adverse events to Melatonin to the specialist, where appropriate.
- Report any adverse events to the MHRA, where appropriate. Report adverse events to the Specialist and MHRA (yellow card reporting scheme).
- To seek advice and/ or help in monitoring the response to and/ or adverse effects to Melatonin treatment.
- Report to and seek advice from the specialist on any aspect of patient care that is of concern to the GP and may affect treatment.
- Stop treatment on advice and/ or correspondence from the specialist
- Stop or adjust treatment if necessary (e.g. side effects) on advice and/ or correspondence from the specialist.
- Continuation without Specialist review is not recommended.
- Continue to support the prescribing of the formulation selected by the specialist where clinical justification has been provided.
- Melatonin treatment should be reviewed by the GP at the 6 month point in between the specialist annual review.

CCG

- To provide feedback to trusts via Trust Medicines Committee.
- To support GPs to make the decision whether or not to accept clinical responsibility for prescribing.
- To support trusts in resolving issues that may arise as a result of shared care.

Patient/Carer

- Contact the GP and community pharmacist to arrange supplies of melatonin in enough time (usually 7 days before re-supply needed) to ensure continuity of treatment.
- Attend appointments.
- Take medication as directed

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

- Report any adverse effects to their GP and/or specialist whilst taking Melatonin.
- Share any concerns in relation to treatment with melatonin.
- Ensure they have a clear understanding of their treatment. Report to the Specialist or GP if they do not have a clear understanding of their treatment.
- Report any changes in disease symptoms to GP and/or specialist whilst taking Melatonin.
- Alert GP and/or specialist of any changes of circumstance which could affect management of disease e.g. plans for pregnancy whilst taking Melatonin.
- If requested, keep a sleep diary to assess the effectiveness of therapy.

Community Pharmacist's Responsibilities

- Order the appropriate product from the wholesaler.
- Inform the patient / carer and GP if there is a supply problem.
- Advise the patient / carer / GP as necessary.

Use in adults

A modified release formulation of melatonin is now licensed and marketed for the short-term treatment of primary insomnia, characterized by poor quality sleep, in patients aged 55 years or over. Its use in this indication is limited to a dose of 2mg at night for no longer than 3 weeks at a time. The use in adults is outside the scope of this guideline and not generally used at the hospital.

CONTACT NUMBERS FOR ADVICE AND SUPPORT

East London Foundation Trust	
Iffah Salim , CAMHS Pharmacist	0207 540 6789
Tsana Simmonds , Lead Pharmacist – Tower Hamlets	0208 223 8014
Andrea Okoloekwe, Lead Pharmacist - Newham	0207 540 4380
Susana FonteloRojano, Lead Pharmacist – City and Hackney	020 8510 8401
Natasha Patel - Luton and Bedfordshire	07940 466861
Clinical Commissioning Groups (CCG)	
CCG Tower Hamlets, Newham and City & Hackney	
Barts Health NHS Trust	
Consultant via switchboard: Registrar on-call out of hours:	020 7377 7000- see top of the document Air call via switchboard
Specialist Pharmacist	020 324 60133

Medicines Information (for drug information related queries)	020 324 60120
CCG Luton and Bedfordshire	
Community paediatrician, Edwin Lobo Centre	01582 700300
Community paediatrician, Union St. clinic	01234 310071

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

Child Development Centre, Hill Rise	01234 310278
Hospital paediatrician, Bedford Hospital	01234 355122

A patient information leaflet suitable for children's parents and carers is attached to the end of the shared care. A copy of this leaflet on Melatonin is also available at:

<http://www.medicinesforchildren.org.uk/search-for-a-leaflet/melatonin-for-sleep-disorders/>

References

1. Weiss M, Wasdell M et al. Sleep hygiene and melatonin treatment for children and adolescents with ADHD and initial insomnia. *Journal of the American Academy of Child and Adolescent Psychiatry* 2006;45:512-519
2. Van der Heijden K, Smits M et al. Effect of melatonin on sleep, behavior, and cognition in ADHD and chronic sleep-onset insomnia. *Journal of the American Academy of Child and Adolescent Psychiatry* 2007;46:233-241
3. Tjon CV et al (2003) Melatonin for treatment of sleeping disorders in children with attention deficit/hyperactivity disorder: a preliminary open label study. *Eur J Pediatr*; 162:554-555
4. LNDG review: Melatonin in paediatric sleep disorders (Sep 2008) <http://www.nelm.nhs.uk/en/NeLM-Area/Evidence/Drug-Specific-Reviews/Melatonin-in-paediatric-sleep-disorders/?query=melatonin&rank=4>
5. Wasdell MB, Jan JE, Bomben MB et al. A randomized, placebo-controlled trial of controlled release melatonin treatment of delayed sleep phase syndrome and impaired sleep maintenance in children with neurodevelopmental disabilities. *J. Pineal Res.* 2008; 44: 57–64
6. Sajith SG, Clarke D (2007) Melatonin and sleep disorders associated with intellectual disability: a clinical review. *Journal of Intellectual Disability Research*; 51(1):2-13
7. Phillips L, Appleton R. Systematic review of melatonin in children with neurodevelopmental disabilities and sleep impairment. *Development Medicine & Child Neurology* 2004;46:771-775
8. Buscemi N et al (2006) Efficacy and safety of exogenous melatonin for secondary sleep disorders and sleep disorders accompanying sleep restriction: meta-analysis. *BMJ*; published online Feb 10th 2006
9. Braam W, Didden R, Smits M, Curfs L (2008) Melatonin treatment in individuals with intellectual disability and chronic insomnia: a randomized placebo-controlled study. *Journal of Intellectual Disability Research*; 52(3):256-264
10. Garstang J, Wallis M (2006) Randomized controlled trial of melatonin for children with autistic spectrum disorders and sleep problems. *Child: care, health and development*; 32(5):585-589
11. Approved by the Bedfordshire and Luton Joint Prescribing Committee April 2013
12. Coppola G., Iervolino G., Mastro Simone M., La Torre G., Ruiu F. & Pascotto A. (2004) Melatonin in wake-sleep disorders in children, adolescents and young adults with mental retardation with or without epilepsy: a double-blind, cross-over, placebo-controlled trial. *Brain and Development* 26, 373–6.
13. Summary of Product Characteristics for Circadin™ (last revised 14/08/2012)
14. <http://www.medicines.org.uk/EMC/medicine/25643/SPC/Circadin/>
15. Great Manchester Interface Prescribing Group. Shared Care Guideline for Melatonin for Sleep Disorder
16. Hampshire Primary Care Trust and Frimley Park Hospital. Shared Care Guideline. Melatonin for sleep disorders in children and adolescents with neurological or behavioural disorders or chronic fatigue syndrome. January 2008.
17. Personal Communication with Penn Pharmaceuticals and Quantum Pharmaceuticals, 20 March 2013.

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

18. BNF for Children 2015/2016
19. Personal Communication, Flynn Pharma (manufacturer of Circadin®), 20 March 2013 (data on file).
20. Drug Tariff (Sept 2016): <http://www.nhsbsa.nhs.uk/PrescriptionServices/4940.aspx>
21. ESUOM2: Sleep disorders in children and young people with attention deficit hyperactivity disorder: melatonin, National Institute for Health and Clinical Excellence, January 2013.
<http://publications.nice.org.uk/esuom2-sleep-disorders-in-children-and-young-people-with-attention-deficit-hyperactivity-disorder-esuom2/key-points-from-the-evidence>
22. LCCG MO: Melatonin Cost Effective Choices Bulletin; Luton CCG (October 2016)
23. Chua, Hui Ming et al. "Dissolution of Intact, Divided and Crushed Circadin Tablets: Prolonged vs. Immediate Release of Melatonin." Ed. Afzal R. Mohammed. *Pharmaceutics* 8.1 (2016): 2. *PMC*. Web. 30 Dec. 2016.

Information Leaflet: Melatonin

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

Melatonin for sleep disorders

This leaflet is about the use of melatonin for particular sleep problems in childhood. It is used to help children who have problems getting to sleep at the start of the night.



This leaflet has been written for parents and carers about how to use this medicine in children. Our information sometimes differs from that provided by the manufacturers, because their information is usually aimed at adult patients. Please read this leaflet carefully. Keep it somewhere safe so that you can read it again.

Name of drug

Melatonin

Common brand: Circadin® (modified-release tablets)

Melatonin is available in a range of branded products.

Why is it important for my child to take this medicine?

Poor sleep can affect your child's physical health, mood, behaviour and development. Melatonin is used to help regulate the sleep pattern of children. Melatonin may help your child to get into a regular sleep pattern.

What is melatonin available as?

- **Modified-release tablets** (Circadin): 2 mg
- **Tablets and capsules** from 0.5 to 5 mg (this is a 'special order' medicine and has to be ordered in advance by your pharmacist)
- **Liquid medicine:** 5 mg per 5 mL (this has to be ordered specially from your pharmacist)

When should I give melatonin?

Melatonin is best given once a day, between half an hour and an hour before your child's agreed bedtime.

Give the medicine at about the same time each day so that this becomes part of your child's daily routine, which will help you to remember.

How much should I give?

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

Your doctor will work out the amount of melatonin (the dose) that is right for *your* child. The dose will be shown on the medicine label.

Your doctor will probably recommend that your child has a low dose to start with. They may then increase the dose until your child's sleep problems have improved. Normally, the dose will not be more than 10 mg per day.

It is important that you follow your doctor's instructions about how much to give.

How should I give melatonin?



Modified-release tablets (Circadin) should be swallowed whole unless your doctor or pharmacist has told you otherwise. Your child should not chew the tablet. Sometimes, your doctor or pharmacist may have told you to crush it – this will make it act faster, but the effect will not last as long.



Other tablets and capsules should be swallowed with a glass of water, milk or juice. You can crush the tablet or open the capsule, and mix the contents with a small amount of soft food such as yogurt, jam or mashed potato. Make sure your child swallows it straight away, without chewing.



Liquid medicine: Measure out the right amount using an oral syringe or medicine spoon. You can get these from your pharmacist. Do not use a kitchen teaspoon as it will not give the right amount.

When should the medicine start working?

Melatonin should start to make your child feel sleepy about half an hour to an hour after taking a dose.

What if my child is sick (vomits)?

- If your child is sick less than 30 minutes after having a dose of melatonin, give them the same dose again.
- If your child is sick more than 30 minutes after having a dose of melatonin, you do not need to give them another dose that night.

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

What if I forget to give it?

If you miss a dose and your child is already asleep, wait until the next day and give the normal dose as usual. If your child is still awake, give them the normal dose.

What if I give too much?

If you think you may have given your child too much melatonin, contact your doctor or local NHS services (111 in parts of England; 0845 4647 in parts of England and Wales; 111 in Scotland).

Take the medicine container or packaging with you, even if it is empty. This will be useful to the doctor. Have the medicine or packaging with you if you telephone for advice.

Are there any possible side-effects?

We use medicines to make our children better, but sometimes they have other effects that we don't want (side-effects).

Side-effects that you must do something about

Rarely, melatonin can cause problems with your child's heart. If your child develops a severe chest pain or has a fast heart rate (they may have a fluttering feeling in their chest or feel their heart beating fast), contact your doctor straight away or take your child to hospital.

If your child seems very unwell in any way that is unusual for them and you are concerned, take them to hospital.

Other side-effects you need to know about

- Your child may feel dizzy or nervous, or may have stomach pain.
- Your child may develop a rash and itch.

If you are concerned about any of these side-effects contact your doctor.

There may, sometimes, be other side-effects that are not listed above. If you notice anything unusual and are concerned, contact your doctor. You can report any suspected side-effects to a UK safety scheme at <http://yellowcard.mhra.gov.uk>.

Can other medicines be given at the same time as melatonin?

- You can give your child medicines that contain paracetamol or ibuprofen, unless your doctor has told you not to.

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

- Check with your doctor or pharmacist **before** giving any other medicines to your child. This includes herbal or complementary medicines.

Is there anything else I need to know about this medicine?

- Treatment with melatonin is usually started by a specialist.
- A specialist may suggest that your child takes just one dose of melatonin before having a CT scan, MRI scan or EEG, when they might be expected to lie still for a while.

General advice about medicines

- Only give this medicine to your child. Never give it to anyone else, even if their condition appears to be the same, as this could do harm.

If you think someone else may have taken the medicine by accident, contact your doctor straight away.

- Make sure that you always have enough medicine. Order a new prescription at least 2 weeks before you will run out.
- Make sure that the medicine you have at home has not reached the 'best before' or 'use by' date on the packaging. Give old medicines to your pharmacist to dispose of.

Where should I keep this medicine?

- Keep the medicine in a cupboard, away from heat and direct sunlight. It does not need to be kept in the fridge.
- Make sure that children cannot see or reach the medicine.
- Keep the medicine in the container it came in.

Who to contact for more information

Your child's doctor, pharmacist or nurse will be able to give you more information about melatonin and about other medicines used to treat sleep disorders.

You can also get useful information from:

The primary source for the information in this leaflet is the British National Formulary for Children. For details on any other sources used for this leaflet, please contact us through our website,

MELATONIN

Insomnia and Sleep Disorders in Children and Adolescents- Shared Care Agreement

Publication date: 05/12/2014 Copyright disclaimer: Version 2, November 2014. © NPPG, RCPCH and WellChild 2011, all rights reserved. Reviewed by: November 2017.

www.medicinesforchildren.org.uk

We take great care to make sure that the information in this leaflet is correct and up-to-date. However, medicines can be used in different ways for different patients. It is important that you ask the advice of your doctor or pharmacist if you are not sure about something. This leaflet is about the use of these medicines in the UK, and may not apply to other countries. The Royal College of Paediatrics and Child Health (RCPCH), the Neonatal and Paediatric Pharmacists Group (NPPG), WellChild and the contributors and editors cannot be held responsible for the accuracy of information, omissions of information, or any actions that may be taken as a consequence of reading this leaflet.