


MANAGEMENT OF SEIZURES IN AN OUT OF HOSPITAL SETTING



Children & Young People

Luton




Parents do not require BLS training

MANAGEMENT OF SEIZURES IN AN OUT OF HOSPITAL SETTING



Children & Young People

Luton



Children & Young People's Seizure Pathway



A seizure is a sudden episode of transient neurological symptoms such as involuntary muscle movements, sensory disturbances and altered consciousness (The American Heritage Science Dictionary).

Obtain blood glucose level at time of seizure or as soon after as possible.
In the hospital setting an ECG should be performed for convulsive non-febrile seizures.

Box 1

Criteria for referral/admission **(where red flags are present, arrange for urgent admission):**

- **Evidence of raised intracranial pressure (blurred vision, bulging fontanelle in infants, persistent headache associated with vomiting)**
- **Drowsy or GCS <15 1 hour post seizure**
- **Signs of sepsis/ meningitis (see fever pathway)**
- **Seizure after a head injury (refer to head injury pathway)**
- Prolonged seizure (lasting more than 5 minutes)
- Recurrent seizure on same day (and these are the first seizures ever experienced by the child). If the patient is known to have cluster seizures in the past, check the care plan or with carers.
- No focus of infection on examination for febrile seizures
- Focal neurological signs (one sided weakness, squint or paraesthesia)
- Age <12 months for febrile seizure
- Age >6 years and 1st febrile seizure
- Age <2 years for non-febrile seizure
- Consider parental anxiety/ social circumstances/safeguarding
- Focal seizures (also known as partial seizures)
 - Motor seizures - 1 side of the body jerking
 - Non-motor seizures- changes in emotions, sensations, thinking or behaviour

This assessment tool was arrived at after careful consideration of the evidence available including but not exclusively NICE Guidelines, EPEN and NHS evidence.

Healthcare professionals are expected to take it fully into account when exercising their clinical judgement. This guidance does not, however, override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

V1.0 - reviewed 24/11/2023